



Medication to treat itching wounds

Class 2 drug

The two components to alleviate itchiness and the component to prevent suppuration penetrate the wound opening to heal the itching wound. This is an oil-based product that will unlikely smart, even when applied, and does not contain steroids.

For Wounds caused by dry skin

For wounds arising after insect bites, and wounds made by scratching off the site touching tight garment elastic MADE IN JAPAN

Indications

Cuts, abrasions, scratches, shoe sores, and skin sterilization/ disinfection

Dosage and Administration

Apply several times a day to the affected area.

(Precautions related to Dosage and Administration)

- (1) Please respect the stipulated application method and dosage.
- (2) Do not touch around the eyes or mouth with the ointment on your hands.
- (3) When used for children, please use under the guidance and supervision of the guardian.
- (4) Keep out of eyes. If by chance the product gets in the eyes, rinse immediately with water or tepid water.
- Furthermore, when symptoms are severe, please receive medical attention from an ophthalmologist.
- (5) For external use only.
- (6) Do not use when the affected part or surrounding area is dirty.

Ingredient, content (per 100g), and action

cetrimide	0.5g	Prevents suppuration by way of sterilizing and disinfecting effect.
lidocaine	2.0g	Profides fast relief of wound itchiness and pain by way of a local anesthesia effect.
diphenhydramine	0.5g	Inhibits wound itchiness by way of an antihistamine action.

Contains micro crystalline wax, glyceryl stearate, sorbitan trioleate, and petroleum jelly as additives.

Precautions

Thigs to consult

1. Before using the product, the following individuals should consult their doctor, pharmacist, or registered sales clerk:

- (1) Individuals receiving medical treatment from a doctor.
- (2) Individuals who have a history of drug-induced allergic reaction.
- (3) Individuals with an extensive area affected
- (4) Individuals with deep wounds, or severe burns

2. After usage, if any of the following symptoms occur, they could be side effects; thus, discontinue usage of the ointment immediately, and please consult your doctor, pharmacist, or registered sales clerk with this document.

Site concerned	Symptoms
Skin	Eruption, redness, itchiness, and swelling

3. If symptoms do not improve after usage for 5 - 6 days, discontinue usage, and please consult your doctor, pharmacist, or registered sales clerk with this document.