



新今治水®

new KONJISUI
(Medicine for toothache)

Medicine for toothache

Class 2 drug

Liquid-type medication for toothache to give fast pain relief by applying directly on the painful tooth area.

MADE IN JAPAN


Indications

For the relief of pain from cavities, floating teeth, and pain of the dental root.

Dosage and Administration

Insert a cotton soaked in the product into the hole of the decayed tooth.

Upon removing the upper cap, a cotton wool ball will be found inside. With clean fingers, make an appropriately sized cotton swab for use.



Soak a cotton swab in an appropriate amount of New Konjisui using the tweezers supplied.



In the event of pain caused by tooth decay with a large cavity, soak a cotton swab with New Konjisui, and push it into the cavity of the decayed tooth.

In the event of pain caused by tooth decay with a small cavity, soak a cotton swab with New Konjisui, and apply it to the surface of the decayed tooth.

Precautions related to Dosage and Administration

- (1) Please respect the stipulated application method.
- (2) Do not apply to areas other than the painful tooth cavity.
- (3) When used for children, please use under the guidance and supervision of the guardian.
- (4) This product is for dental use only. Do not use for ophthalmological or other uses.
- (5) This product will not harm the hard quality of the tooth (enamel and dentine quality); however, if excessive product spreads to areas other than the tooth, taste will temporarily change, and so please wipe of such as with a gauze.
- (6) If the product gets on the face or around the mouth by mistake, rinse immediately with water and wipe the product off.

※This product cannot cure the cause of toothache such as tooth decay, and so it is recommended to receive fundamental treatment from a dentist without delay.

Ingredient, content (per 100g), and action

Ingredients	Content	Action
clove oil	0.3g	Pain relief
phenol	5.0g	Pain relief and bactericidal action.

camphor	10.0g	Pain relief
Cinnamon oil	2.5g	
l-menthol	0.1g	
diphenhydramine hydrochloride	2.0g	Inhibits inflammation.
dibucaine hydrochloride	0.1g	Pain relief
p-butylaminobenzoyldiethylaminoethanol hydrochloride	0.03g	
gardenia tincture	15.07g	Inhibits inflammation.

Contains propylene glycol and ethanol as additive.

Precautions

Thigs to avoid

(if not respected, the present symptoms will exacerbate, and side effects will readily develop)

Do not use the product in the following areas:

Gums, and lips

Thigs to consult

- Before using the product, the following individuals should consult their doctor, dentist, pharmacist, or registered sales clerk:
 - Individuals receiving medical treatment from a doctor or dentist.
 - Individuals who have a history of drug-induced allergic symptoms.
- After usage, if any of the following symptoms occur, they could be side effects, and thus, discontinue usage of the product immediately, and please consult your doctor, pharmacist, or registered sales clerk with this document.

Site concerned	Symptoms
Skin	Eruption, redness, and itchiness

- If symptoms do not improve after usage for 5 - 6 times, discontinue usage, and please consult your doctor, dentist, pharmacist, or registered sales clerk with this document.